

## **Certificate of Analysis**

PRODUCT: FlyCut® PstI

CAT. NO. AMOUNT JP101 2\*250 units

**LOT NUMBER**P20528

SHELF LIFE
Two years

STORAGE CONDITIONS SHIPPING CONDITIONS

at -20 °C for two years Dry ice(-70 °C)

### **DESCRIPTION**

*FlyCut*<sup>®</sup> PstI is expressed and purified from *E.coli* that carries the recombinant PstI gene. The molecular weight is 38.3 kDa, with the recognition site at CTGCA^G. The reaction is conducted for 5-15 minutes at 37°C, and heat-inactivated at 80°C for 20 minutes. This enzyme is not sensitive to dam, dcm or mammalian CpG methylation.

### **Kit Contents**

Component	JP101-01	JP101-02
FlyCut® PstI	5000 units	2×5000 units
10×FlyCut® Buffer	2×1 ml	3×1 ml
10×DNA Loading Buffer	1 ml	1 ml



# **Certificate of Analysis**

### QUALITY CONTROL DATA

Assay	Standard	Result
Appearance	The liquid should be clear and free of precipitates or floccules.	Met standards
16 hours Incubation	In a 50 µl reaction system, 10 units of enzyme are incubated with 1 µg DNA for 16 hours. The result is comparable with that of 1-hour incubation with 1 unit of enzyme.	Met standards
Ligation and Recutting	Digested with a 10-fold excess of endonuclease, more than 95% of the DNA fragments can be ligated with T4 DNA ligase at 25°C, and 95% of the ligated fragments can be recut.	Met standards
Terminal Integrity	The linearized plasmid containing the LacZ $\alpha$ gene (the only site) is religated and transformed on a plate coated with X-gal/IPTG. The successful expression of $\beta$ -galactosidase is the criterion for detecting the integrity of the gene after cloning. Blue colonies indicate the integrity of the gene, while white colonies which indicate the gene is damaged must be less than 3% of the total number of colonies.	Met standards
Endonuclease Activity	In a 50 $\mu$ l reaction system, incubation of 15 units of enzyme with 1 $\mu$ g of pBR322 DNA at 37°C for 4 hours results in no more than 10% from RFI to RFII.	Met standards

## **Notice to Purchaser**



All products are for research use only. Caution: Not intended for human or animal diagnostic or therapeutic uses. If you have any further questions about this Certificate of Analysis, please contact us at +86-10-57815027 or +86-400-898-0321.

TransGen Biotech products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products or to provide a service to third parties without prior written approval of TransGen Biotech Co., Ltd.

TransGen Biotech Co., Ltd. Building No.4, Zhongguancun Dongsheng International Science Park, No. 1 North Yongtaizhuang Road, Haidian District, Beijing, China

Postcode: 100192

Tel: +86-10-57815027 Fax: +86-10-57815000

Customer Service: +86-400-898-0321 Web site: www.transgenbiotech.com E-mail: trans@transgen.com.cn info@transgenbiotech.com

Manager, Quality Control Department

Heng Zheng

31/05/2021